If it's going into your patient's body, **quality matters** more than ever

Every time you flush a patient's line,
you are injecting saline directly into their bloodstream

Have you carefully considered the quality of your prefilled flush syringe?

Not all prefilled flush syringes are created equal[1]

Quality defects and bacterial contamination in prefilled flush syringes have **seriously harmed patients**[2,3,4]:

Catheter-related infections[1]

Spinal meningitis[1]

Permanent brain damage[1]

11 deaths, including a 2-year-old child, were reported in the U.S. from the use of contaminated syringes from 2008 to 2017

With **BD PosiFlush™ Syringes**, you can be confident in the **quality** of *every flush, every time.*

Because each **BD PosiFlush™ Syringe** is created with three criteria:

SAFETY | **EXPERIENCE** | **RELIABILITY**

Be confident it's a **BD PosiFlush™ Syringe**

If it's going into your patient's body,

SAFETY
matters more than ever

Be confident it's a BD PosiFlush™ Syringe

Touch-free manufacturing

ensuring the first person to touch the syringe is **the clinician**[5]

CE-certified Class-III Medical Device

unlike other prefilled saline syringes classified as Class-IIa CE-certified devices, BD PosiFlush™ Syringes are classified as Class-III Medical Devices.

Robust multi-step inspection process

to meet the highest quality standards and safety for our customers and patients[5]

All steps controlled

in the manufacturing process, enabling us to manage the quality process end-to-end [5]

Be confident it's a BD PosiFlush™ Syringe

If it's going into your patient's body,

EXPERIENCE
matters more than ever

Be confident it's a BD PosiFlush™ Syringe

20+ years
with BD PosiFlush™ Syringes in the market as a partner, supporting the needs of patients globally[6]

18 billion+
BD PosiFlush™ Syringes produced through 2022[5]

13,000+ customers
around the world use and trust BD PosiFlush™ Syringes[7]

Be confident it's a
BD PosiFlush™ Syringe

If it's going into your patient's body,

RELIABILITY
matters more than ever

Be confident it's a BD PosiFlush™ Syringe

>25% production increase on **BD PosiFlush™ SP Syringe** (2020-2022)

delivering to our customers when market changes necessitated additional support [8]

3 YEARS

3-year shelf life

enabling facilities to plan ahead and manage inventory [9]

Following 6 health organizations'

best practice for flushing: INS, RNC, NICE, epic 3, INICC, ISMP, NHS Standards [10-16]

*INS: Infusion Nurses Society; RCN: Royal College of Nursing; NICE: National Institute for Health and Care Excellence; INICC: International Nosocomial Infection Control Consortium; ISMP: Institute for Safe Medication Practice. NHS: National Health Service.

We are commited to meet our customers' needs by ensuring responsive and timely delivery

supported by 3 manufacturing plants and our global network of distribution centres

Note: Boxes of images and captions shown below will only appear on hover in final Infogram

Drogheda, Ireland

Columbus, Nebraska, USA

Fraga, Spain

ISO-compliant

with relevant standards for quality assurance and environmental best practices

20+ published clinical studies [12–14]

more than any other product in the prefilled flush category*

*Based on a Pubmed literature search of key saline flush competitors as of July 2023.

Be confident it's a **BD PosiFlush™ Syringe**

Because so much is on the line

Learn more about how you can help ensure delivery of safety, experience and reliability

Be confident, it's a BD PosiFlush™ Syringe

[CTA: Link to product page or lead-gen form]

References:

1. Massoomi F, Arruda J. A Closer Look at Prefilled Flush Syringe Safety. Pharmacy Practice News. February 15, 2018. Accessed on December 18, 2023 at https://www.pharmacypracticenews.com/Clinical/Article/07-14/A-Closer-Look-at-Prefilled-Flush-Syringe-Safety/46801
2. Jewett C, Nguyen D. Timeline: How Contaminated Syringes Got From Factory to Patients. ProPublica. June 6, 2009. Accessed on May 2, 2023, at https://www.propublica.org/article/timeline-am2pat.
3. U.S. Food and Drug Administration. Cardinal Health Issues Nationwide Recall of Select Monoject Flush Prefilled Syringes. Aug 2021. Accessed on May 2, 2023,
4. Center for Disease Control and Prevention (CDC).Update: Delayed Onset Pseudomonas fluorescens Bloodstream Infections After Exposure to Contaminated Heparin Flush. Web September 7, 2006. Accessed on November 13th, 2023.
5. Gorski LA, Hadaway L, Hagle ME, et al. Infusion Therapy Standards of Practice, 8th Edition. J Infus Nurs. 2021;44(1S,1):S1-S224. doi: 10.1097/NAN.0000000000000396
6. Denton A, Bodenham A, Conquest A, et al. Royal College of Nursing. Standards for Infusion Therapy , 4th Edition. https://www.rcn.org.uk/-/media/royal-college-of-nursing/documents/publications/obselete/005704.pdf?la=en. Published December 12, 2016. Accessed June 2, 2021.
7. Loveday HP, Wilson JA, Pratt RJ, et al. epic3: National Evidence-Based Guidelines for Preventing Healthcare-Associated Infections in NHS Hospitals in England. J Hosp Infect . 2014;86(1):S1-S70. doi: 10.1016/S0195-6701(13)60012-2
8. Rosenthal VD, Kanj SS, Desse J, et al. Bundle of the International Nosocomial Infection Control Consortium (INICC) to Prevent Central and Peripheral Line-Related Bloodstream Infections. The International Nosocomial Infection Control Consortium (INICC). Published February 11, 2017. Accessed October 20, 2020. doi: 10.13140/RG.2.2.19556.99200
9. National Health Service (NHS) National Patient Safety Agency. NPSA Alert - Promoting safer use of injectable medicines (NPSA 20). https://www.sps.nhs.uk/articles/npsa-alert-promoting-safer-use-of-injectionmedicinesnpsa-20-2007/. Published March 28, 2007. Updated August 22, 2018. Accessed October 4, 2019.
10. The Institute for Safe Medication Practices (ISMP). ISMP Safe Practice Guidelines for Adult IV Push Medications. ISMP . http://www.ismp.org/sites/default/files/attachments/2017-11/ISMP97-Guidelines-071415-3.%20FINAL.pdf. Published July 14, 2015. Accessed October 20, 2020.
11. Joint Commission The Joint Commission. National Patient Safety Goals Effective July 2020 for the Hospital Program. The Joint Commission. http://www.jointcommission.org/-/media/tjc/documents/standards/national-patient-safety-goals/2020/npsg_chapter_hap_jul2020.pdf. Published March 26, 2020. Accessed October 20, 2020.
12. Keogh S, Marsh N, Higgins N, Davies K, Rickard C. A time and motion study of peripheral venous catheter flushing practice using manually prepared and pre-filled flush syringes. J Infus Nurs . 2014;37(2):96-101. doi: 10.1097/NAN.0000000000000024
13. Saliba P, Cuervo G, Hornero A, et al. The impact of flushing with pre-filled saline syringes on the incidence of peripheral venous catheter failure: A quasi-experimental study. J Vasc Access. 2019:21(4); 490-496. doi: 10.1177/1129729819888423
14. Bertoglio S, et al. Improving outcomes of short peripheral vascular access in oncology and chemotherapy administration. J Vasc Access. 2017;18(2):89-96. doi:10.5301/jva.5000668

CE 0050

BD–Europe, Eysins, Switzerland, 1262

BD, the BD Logo and PosiFlush are trademarks of Becton, Dickinson and Company or its affiliates. All other trademarks are the property of their respective owners.
© 2023. All rights reserved. BD 92921 (12/2023)

BD